


## Determination of Pesticides in Grape Juices by QuEChERS and Liquid Chromatography-Tandem Mass Spectrometry

Deise F. Souza,<sup>a,b</sup> Edson L. Souza<sup>a</sup> and Endler M. Borges<sup>\*,a,c</sup>


<sup>a</sup>Núcleo Biotecnológico, Universidade do Oeste de Santa Catarina, UNOESC, Rua Paese, 198, Bairro Universitário-Bloco K, 89560-000 Videira-SC, Brazil

<sup>b</sup>Laboratório da Tecnologia de Bebidas, LATEB, Núcleo de Metrologia Gestão e Processo Produtivo, Sistema FIESC-SENAI/SC Videira, Rua Julio Pretti, 270, 89570-000 Pinheiro Preto-SC, Brazil


<sup>c</sup>Departamento de Química, Universidade Regional de Blumenau, FURB, Campus 1, Rua Antônio da Veiga, 140, Victor Konder, 89012-900 Blumenau-SC, Brazil




**Figure S1.** Matrix effect. Black line TIC of a fortified sample at 2 mg L<sup>-1</sup>. Red line TIC of standard solution at 2 mg L<sup>-1</sup>.



**Figure S2.** LC-MS/MS analysis of azoxystrobin in grape juice. (a) TIC; (b) qualifier  $404 \rightarrow 372$ ; (c) quantifier transitions  $404 \rightarrow 344$ .



**Figure S3.** LC-MS/MS analysis of metalaxyl in grape juice. (a) TIC; (b) qualifier  $280 \rightarrow 220$ ; (c) quantifier transitions  $280 \rightarrow 160$ .



**Figure S4.** LC-MS/MS analysis of tebuconazole in grape juice. (a) TIC; (b) qualifier  $308 \rightarrow 70$ ; (c) quantifier transitions  $308 \rightarrow 124$ .