

Catalyst- and Solvent-free Synthesis of Imidazo[1,2-a]pyridines

Dong-Jian Zhu,^a Jiu-Xi Chen,^{*a} Miao-Chang Liu,^a Jin-Chang Ding,^{a,b} and Hua-Yue Wu^{*,a}

^aCollege of Chemistry and Materials Engineering, Wenzhou University, Wenzhou, 325027, China

^bWenzhou Vocational and Technical College, Wenzhou, 325035, P. R. China




Figure S1. ¹H NMR of 3a (300 MHz, CDCl₃) and ¹³C NMR of 3a (75 MHz, CDCl₃).

*e-mail: jiuxichen@wzu.edu.cn; huayuewu@wzu.edu.cn




Figure S2. ¹H NMR of **3b** (300 MHz, CDCl₃) and ¹³C NMR of **3b** (75 MHz, CDCl₃).




Figure S3. ¹H NMR of **3c** (300 MHz, CDCl₃) and ¹³C NMR of **3c** (75 MHz, CDCl₃).





Figure S4. ¹H NMR of **3d** (300 MHz, CDCl₃) and ¹³C NMR of **3d** (75 MHz, CDCl₃).






Figure S6. ^1H NMR of **3f** (300 MHz, CDCl_3) and ^{13}C NMR of **3f** (75 MHz, CDCl_3).




Figure S7. ¹H NMR of **3g** (300 MHz, CDCl₃) and ¹³C NMR of **3g** (75 MHz, CDCl₃).




Figure S8. ¹H NMR of **3h** (300 MHz, CDCl₃) and ¹³C NMR of **3h** (75 MHz, CDCl₃).




Figure S9. ¹H NMR of **3i** (300 MHz, CDCl₃) and ¹³C NMR of **3i** (75 MHz, CDCl₃).




Figure S10. ¹H NMR of **3j** (300 MHz, CDCl₃) and ¹³C NMR of **3j** (75 MHz, CDCl₃).




Figure S11. ¹H NMR of **3k** (300 MHz, CDCl₃) and ¹³C NMR of **3k** (75 MHz, CDCl₃).




Figure S12. ¹H NMR of **31** (300 MHz, CDCl₃) and ¹³C NMR of **31** (75 MHz, CDCl₃).




Figure S13. ¹H NMR of **3m** (300 MHz, CDCl₃) and ¹³C NMR of **3m** (75 MHz, CDCl₃).