

A New and Concise Strategy to the Enantioselective Synthesis of
(S)-2-Amino-4-Oxo-4-(Pyridine-2-yl) Butanoic Acid from Aspartic Acid

Evanoel Crizanto de Lima,^a Paulo G. de Lima,^b Carolina C. de Souza,^a
Marta C. L. S. Maior,^a Ayres G. Dias^c and Paulo R. R. Costa^{*a}

^aNúcleo de Pesquisas de Produtos Naturais, Centro de Ciências da Saúde, Universidade Federal do Rio de Janeiro, 21941-540 Rio de Janeiro-RJ, Brazil

^bInstituto de Química, Universidade Federal do Rio de Janeiro, 21941-909 Rio de Janeiro-RJ, Brazil

^cInstituto de Química, Universidade do Estado do Rio de Janeiro, 22250-040 Rio de Janeiro-RJ, Brazil




Figure S1. ¹H NMR Spectrum of compound 6.




Figure S2. ¹H NMR Spectrum of compound 7.




Figure S3. ¹H NMR Spectrum of compound 13.




Figure S4. ^{13}C NMR Spectrum of compound 13.




Figure S5. Low resolution mass spectrometry of compound 13.




Figure S6. ¹H NMR Spectrum of compound 5.




Figure S7. ¹³C NMR Spectrum of compound 5.




Figure S8. ¹H NMR Spectrum of compound 8.




Figure S9. Low resolution mass spectrometry of compound 8.




Figure S10. ¹H NMR Spectrum of compound 9.