

1,3-Thiazolium-5-thiolates Mesoionic Compounds: Semiempirical Evaluation of their First Static Hyperpolarizabilities and Synthesis of New Examples

Bruno F. Lyra,^{a,b} Soraya A. de Moraes,^a Gerd B. Rocha,^{*,a} Joseph Miller,^{b,c,†}
Gustavo L. C. Moura,^c Alfredo M. Simas,^c Cloviseppe^d and Petrônio F. de Athayde-Filho^{a,b}

^aDepartamento de Química, Universidade Federal da Paraíba, 58059-900 João Pessoa-PB, Brazil

^bLaboratório de Tecnologia Farmacêutica, Universidade Federal da Paraíba, CP 5009,
58051-970 João Pessoa-PB, Brazil

^cDepartamento de Química Fundamental, Universidade Federal de Pernambuco,
50670-901 Recife-PE, Brazil

^dDepartamento de Química, Universidade Federal de Santa Maria,
97105-900 Santa Maria-RS, Brazil




Figure S1. ¹³C (APT) NMR spectrum for MIC-14 (CDCl₃, 50 MHz).

*e-mail: gbr@quimica.ufpb.br

† in memoriam




Figure S2. Expanded version of ¹³C (APT) NMR spectrum for MIC-14 (CDCl₃, 50 MHz).




Figure S3. ¹H NMR spectrum for MIC-14 (CDCl₃, 200 MHz).




Figure S4. Expanded version of ^1H NMR spectrum for MIC-14 (CDCl_3 , 200 MHz).




Figure S5. ^{13}C (APT) NMR spectrum for MIC-16 (CDCl_3 , 50 MHz).




Figure S6. Expanded version of ^{13}C (APT) NMR spectrum for MIC-16 (CDCl_3 , 50 MHz).




Figure S7. ^1H NMR spectrum for MIC-16 (CDCl_3 , 200 MHz).




Figure S8. Expanded version of ¹H NMR spectrum for MIC-16 (CDCl₃, 200 MHz).