## Two Novel Eremophilane Sesquiterpenes from an Endophytic Xylariaceous Fungus Isolated from Leaves of *Cupressus lusitanica*

Luciana S. Amaral and Edson Rodrigues-Filho\*

Departamento de Química, Universidade Federal de São Carlos, CP 676, 13565-905 São Carlos-SP, Brazil



Figure S1. <sup>1</sup>H NMR spectrum of compound 1 (400 MHz, CDCl<sub>3</sub>).

\*e-mail: edson@dq.ufscar.br



Figure S2. <sup>13</sup>C NMR spectrum of compound 1 (100 MHz, CDCl<sub>3</sub>).



Figure S3. DEPT 135 spectrum of compound 1 (100 MHz, CDCl<sub>3</sub>).



Figure S4. <sup>1</sup>H-<sup>1</sup>H COSY NMR correlation spectroscopy 2D NMR spectrum of compound 1 (400 MHz, CDCl<sub>3</sub>).



Figure S5. <sup>1</sup>H-<sup>13</sup>C HSQC 2D NMR correlation spectroscopy of compound 1 (400 MHz/100 MHz, CDCl3).



Figure S6. <sup>1</sup>H-<sup>13</sup>C HMBC 2D NMR correlation spectroscopy of compound 1 (400 MHz/100 MHz, CDCl<sub>3</sub>).



Figure S7. nOe spectra for H-8, H-3, H-6b and H-6a of compound 1 (400 MHz, CDCl<sub>3</sub>).



Figure S8. <sup>1</sup>H NMR spectrum of compound 2 (400 MHz, CDCl<sub>3</sub>).



Figure S9. <sup>1</sup>H-<sup>1</sup>H COSY NMR correlation spectroscopy 2D NMR spectrum of compound 2 (400 MHz, CDCl<sub>3</sub>).



Figure S10. <sup>1</sup>H-<sup>13</sup>C HSQC 2D NMR correlation spectroscopy of compound 2 (400 MHz/100 MHz, CDCl3).



Figure S11. <sup>1</sup>H-<sup>13</sup>C HMBC 2D NMR correlation spectroscopy of compound 2 (400 MHz/100 MHz, CDCl<sub>3</sub>).