

Supplementary Information

Bioactive Phenylpropanoids from *Daphne feddei*

Yanling Lu,^{a,b} Xuesen Li,^{a,b} Huaixue Mu,^a Haitao Huang,^b Gan-Peng Li^{*a} and Qiufen Hu^{*a}

^aKey Laboratory of Ethnic Medicine Resource Chemistry (Yunnan University of Nationalities),
State Ethnic Affairs Commission & Ministry of Education, Kunming 650031, P. R. China.

^bKey Laboratory of Tobacco Chemistry of Yunnan Province, Yunnan Academy of Tobacco Science,
Kunming 650106, P. R. China




Figure S1. DEPT and ¹³C NMR (C_5D_5N , 125 MHz) spectra of feddeiketone A (1).




Figure S2. ¹H NMR ($\text{C}_5\text{D}_5\text{N}$, 500 MHz) spectra of feddeiketone A (**1**).






Figure S3. HSQC spectra of feddeiketone A (**1**).

**Figure S4.** HMBC spectra of feddeiketone A (**1**).**Figure S5.** HRESIMS of feddeiketone A (**1**).




Figure S6. DEPT and ¹³C NMR ($\text{C}_5\text{D}_5\text{N}$, 125 MHz) spectra of feddeiketone B (2).




Figure S7. ¹H NMR ($\text{C}_5\text{D}_5\text{N}$, 500 MHz) spectra of feddeiketone B (2).




Figure S8. DEPT and ¹³C NMR ($\text{C}_5\text{D}_5\text{N}$, 125 MHz) of feddeiketone C (3).




Figure S9. ¹H NMR ($\text{C}_5\text{D}_5\text{N}$, 500 MHz) of feddeiketone C (3).