

Supplementary Information

Novel Bioactive Dibenzocyclooctadiene Lignans from *Schisandra neglecta*

Xue-Mei Gao,^{a,b} De-Yun Niu,^a Chun-Yang Meng,^a Fu-Quan Yao,^a Bin Zhou,^a
Rui-Rui Wang,^c Liu-Meng Yang,^c Yong-Tang Zheng,^c Qiu-Fen Hu,^{*,a}
Han-Dong Sun^b and Wei-Lie Xiao^{*,b}

^aKey Laboratory of Ethnic Medicine Resource Chemistry (Yunnan University of Nationalities),
State Ethnic Affairs Commission & Ministry of Education, Kunming 650031, P. R. China

^bState Key Laboratory of Phytochemistry and Plant Resources in West China,
Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650204, P. R. China

^cKunming Institute of Zoology, Chinese Academy of Sciences, Kunming 650223, P. R. China




Figure S1. ¹³C NMR (C₅D₅N, 125 MHz) and DEPT of neglectalignan A (1).




Figure S2. ^1H NMR ($\text{C}_5\text{D}_5\text{N}$, 500 MHz) of neglectalignan A (**1**).




Figure S3. HSQC of neglectalignan A (**1**).




Figure S6. HRMS of neglectalignan A (1).




Figure S7. CD (MeOH) of neglectalignan A (1).




Figure S8. ^{13}C NMR ($\text{C}_5\text{D}_5\text{N}$, 125 MHz) and DEPT of neglectaligan B (2).




Figure S9. ^1H NMR ($\text{C}_5\text{D}_5\text{N}$, 500 MHz) of neglectaligan B (2).




Figure S10. ^{13}C NMR (CDCl₃, 125 MHz) of neglectalignan C (3).




Figure S11. ^1H NMR (CDCl₃, 500 MHz) of neglectalignan C (3).




Figure S12. ^{13}C NMR ($\text{C}_3\text{D}_5\text{N}$, 125 MHz) of neglectalignan D (4).




Figure S13. ^1H NMR ($\text{C}_3\text{D}_5\text{N}$, 500 MHz) of neglectalignan D (4).