

# Supplementary Information

## A Novel, Donor-Active Solvent-Assisted Liquid-Phase Microextraction Procedure for Spectrometric Determination of Zinc

Livia Kocúrová,<sup>\*a</sup> Ioseph S. Balogh,<sup>b</sup> Martina Fatľová,<sup>a</sup> Yaroslav Bazel',<sup>a</sup> András Simon,<sup>c</sup>  
Rastislav Serbin,<sup>a</sup> Miroslav Badida,<sup>d</sup> Radoslav Rusnák<sup>d</sup> and Vasil Andruch<sup>a</sup>

<sup>a</sup>Department of Analytical Chemistry, University of P. J. Šafárik, SK-04154 Košice, Slovakia

<sup>b</sup>Department of Chemistry, College of Nyíregyháza, HU-4400 Nyíregyháza, Hungary

<sup>c</sup>Department of General and Analytical Chemistry, Budapest University of Technology and Economics, HU-1111 Budapest, Hungary

<sup>d</sup>Department of Environmentalistics, Faculty of Mechanical Engineering, Technical University of Košice, Park Komenského 5, SK-04154 Košice, Slovakia




Figure S1. <sup>1</sup>H NMR spectrum (500 MHz, methanol-d<sub>4</sub>) of DTVTI (a) and magnification of the <sup>1</sup>H NMR spectrum between 6.00 and 8.40 ppm.

\*e-mail: liviamonika.kocurova@gmail.com