

Supplementary Information

Caesalpinioflavone, a New Cytotoxic Biflavonoid Isolated from *Caesalpinia pluviosa* var. *peltophoroides*

João L. B. Zanin,^a Murilo Massoni,^a Marcelo H. dos Santos,^a Giovana C. de Freitas,^b
Evandro L. O. Niero,^c Renata R. Schefer,^b João H. G. Lago,^d Marisa Ionta^e and
Marisi G. Soares*^a

^aInstituto de Química and ^eInstituto de Ciências Biomédicas, Universidade Federal de Alfenas,
37130-000 Alfenas-MG, Brazil

^bInstituto de Química and ^cInstituto de Ciências Biomédicas, Universidade de São Paulo,
05508-000 São Paulo-SP, Brazil

^dInstituto de Ciências Ambientais, Químicas e Farmacêuticas, Universidade Federal de São Paulo,
09920-000 São Paulo-SP, Brazil




Figure S1. HRESIMS for caesalpinioflavone in ESI negative mode.




Figure S2. IR spectrum (KBr) of caesalpinioflavone.




Figure S3. ¹H NMR spectrum (500 MHz, CD₃OD) of caesalpinioflavone.




Figure S4. ¹³C NMR spectrum (125 MHz, CD₃OD) of caesalpinioflavone.




Figure S5. COSY90 spectrum (500 MHz, CD₃OD) of caesalpinioflavone.




Figure S6. HSQC spectrum (500 and 125 MHz, CD₃OD) of caesalpinioflavone.




Figure S7. HMBC spectrum (125 MHz, CD₃OD) of caesalpinioflavone.