

Artigo

First-time Isolation of Flavonoids and Cytotoxic Potential of the Amazonian Shrub *Ptychopetalum olacoides* Benth

Dutra, K. D. B.;[#] Macedo, A. L.;[#] Montenegro, R. C.; Jimenez, P. C.; Castro, R. N.; Epifanio, R. A.;[†] Vasconcelos, T. R. A.; Valverde, A. L.*

Rev. Virtual Quim., 2017, XX (XX), no prelo. Data de publicação na Web: XX de XXXX de XXXX

<http://rvq.sbq.org.br>

Primeiro Isolamento de Flavonoides e Potencial Citotóxico do Arbusto Amazônico *Ptychopetalum olacoides* Benth

Resumo: No presente estudo, três flavonoides, 3-*O*-metilquercetina (**1**), 3,4'-*O*-dimetilquercetina (**2**) e 3,7-*O*-dimetilquercetina (**3**), foram isolados e caracterizados pela primeira vez a partir do extrato metanólico da espécie *Ptychopetalum olacoides* Benth. As estruturas das substâncias foram elucidadas por métodos espectroscópicos (1D-, 2D-RMN, EM e UV) e confirmadas por comparação com a literatura. A atividade citotóxica do extrato bruto foi avaliada *in vitro* contra três linhagens de células humanas cancerígenas. Foi observada atividade moderada (CI₅₀ = 45.16 µg/mL) contra a linhagem de câncer de mama (MCF-7) e, além disso, o extrato bruto não foi citotóxico contra a linhagem de fibroblastos humanos não cancerígenos (MRC-5).

Palavras-chave: Antitumoral; câncer; flavonoides; Olacaceae; *P. olacoides*.

Abstract

In the present study, three flavonoids, 3-*O*-methylquercetin (**1**), 3,4'-*O*-dimethylquercetin (**2**) and 3,7-*O*-dimethylquercetin (**3**) were isolated and characterized for the first time from a methanol extract obtained from the species *Ptychopetalum olacoides*. The structures of compounds were identified by spectroscopic methods (1D-, 2D-NMR, MS and UV) and confirmed by comparison with the respective literature data. The cytotoxic effect of crude extract was evaluated *in vitro* against three human cancer cell lines. The results showed a mild cytotoxic activity (IC₅₀ = 45.16 µg/mL) against breast cancer (MCF-7). However, crude extract did not exhibit any cytotoxic effect against normal cell human fibroblast (MRC-5).

Keywords: Antitumor; cancer; flavonoids; Olacaceae; *P. olacoides*.

* Universidade Federal Fluminense, Instituto de Química, Departamento de Química Orgânica, Campus do Valonguinho, 24020-141, Niterói-RJ, Brazil.

[#] Shared first author position.

[†] *in memoriam*

✉ alessandravalverde@id.uff.br

DOI:

First-time Isolation of Flavonoids and Cytotoxic Potential of the Amazonian Shrub *Ptychopetalum olacoides* Benth

Karen D. B. Dutra,^{a,#} Arthur L. Macedo,^{a,#} Raquel C. Montenegro,^b Paula C. Jimenez,^c Rosane N. Castro,^d Rosângela de A. Epifanio,^{a,+} Thatyana R. A. Vasconcelos,^a Alessandra L. Valverde,^{a,*}

^a Universidade Federal Fluminense, Instituto de Química, Departamento de Química Orgânica, Campus do Valonguinho, CEP 24020-141, Niterói-RJ, Brazil.

^b Universidade Federal do Pará, Instituto de Ciências Biológicas, CEP 66075-110, Belém-PA, Brazil.

^c Universidade Federal de São Paulo, Departamento de Ciências do Mar, CEP 11070-100, Santos-SP, Brazil.

^d Universidade Federal Rural do Rio de Janeiro, Departamento de Química, CEP 23897-000, Seropédica-RJ, Brazil.

* alessandravalverde@id.uff.br

Recebido em 23 de março de 2017. Aceito para publicação em 27 de novembro de 2017

1. Introduction
2. Materials and methods
 - 2.1. General procedures
 - 2.2. Plant material
 - 2.3. Extraction and isolation
 - 2.4. Identification
 - 2.5. Cytotoxic activity assay
3. Results and discussion
4. Conclusion

1. Introduction

Ptychopetalum olacoides Benth. (Olacaceae) is a shrub or small tree widely known in Brazil as “muirapuama”, “marapuama”, “marapuana” and “muiratã”.¹ This is an endemic species to the Amazon rainforest and specially distributed in the

north region of the country in Amazonas, Amapá and Pará states.² Preparations with the stems of *P. olacoides* have been used to treat “nervous weakness”, fatigue, depression symptoms, tremor disorders, and sexual dysfunction.³ The fluid root extract of this plant has been employed in phytotherapeutic formulations as Catuama®, a general tonic widely used in some regions

of Brazil.⁴ However, there are few data related to the phytochemical profile of this species. Montrucchio and co-workers previously reported the isolation of saturated fatty acids (stearic and palmitic acids), methylxanthine caffeine, triterpenoid lupeol and steroid β -sitosterol.⁵ Other studies described the isolation of clerodane diterpenoids,⁶ benzoic acid derivatives such as vanillic and protocatechuic acids, and methylxanthine theobromine.⁷ Despite validation of the total flavonoids content from *P. olacoides*,⁸ there are no reports concerning isolation and characterization of flavonoids in this genus.

2. Materials and methods

2.1. General procedures

TLC was performed on plates pre-coated with silica gel 60 F₂₅₄ (Merck, Germany). Preparative HPLC was performed on a Phenomenex C18 (30 cm x 10 mm x 5 μ m, Torrance, Canada) equipped with a Shimadzu LC-10AS pump and a SPD10A UV/Vis detector (Shimadzu, Kyoto, Japan). The UV spectra were recorded on a JASCO V-370 Bio spectrophotometer (Tokyo, Japan). The NMR spectra were measured on Varian VNMRS 500 MHz spectrometer for ¹H and 125 MHz for ¹³C (Palo Alto, USA), and chemical shifts were reported in ppm downfield from TMS. The MS data were recorded on a Flexar SQ 300 LC/MS system (PerkinElmer, Shelton, CT, USA) using an analytical C18 column (PerkinElmer, 150 mm x 4.6 mm, 3 μ m). A micro-splitter valve was used to send 45% of the flow to the mass spectrometer. The quadrupole mass spectrometer equipped with electrospray ionization (ESI-MS) was operated under positive ion mode. The MS parameters were set at 12 L/min for drying gas flow, 80 psi for nebulizer pressure and 300 °C for drying gas temperature. Column chromatography was carried out on Sephadex LH-20.

2.2. Plant material

The powdered wood/bark of *P. olacoides* was acquired from Santosflora Herbs Ltda. (CNPJ: 51569309/0001-38, IBAMA registration No. 35867 and ANVISA's authorization No. 6.02.671-1) in June 2013. The species was collected in February 2013, lot code MARPP01/0213 and validity period from 02.04.2013 to 02.04.2016.

2.3. Extraction and isolation

The bark and wood powder of *P. olacoides* (500 g) was extracted with a solvent gradient of increasing polarity under sonication using *n*-hexane, ethyl acetate and methanol. After extraction and removal of the solvent under vacuum in a rotatory evaporator, a dark residue was obtained (4.10 g) from the methanol extract (ME). The ME fraction (2.6 g) was restructured in methanol and filtered using Fisherbrand nylon 0.2 μ m filter to obtain a particle-free extract. The extract was then chromatographed on lipophilic Sephadex LH-20 (25-100 μ m) and eluted with MeOH,⁹ resulting mainly in 12 fractions (MEF1-12). MEF8 (20 mg) was subsequently purified by preparative RP-HPLC [mobile phase: H₂O/AcOH (99:1) (solvent A) and MeOH (solvent B) at a constant flow rate of 5 mL/min, using 65% solvent B and 35% solvent A, detection at 340 nm] resulting in substances **1** (1.0 mg), **2** (1.0 mg), and **3** (1.0 mg).

2.4. Identification

The NMR spectra of compounds **1-3** were acquired with ¹H, COSY, HSQC and HMBC techniques. The MS data were obtained by LC/MS analysis of MEF8 using as mobile phase a gradient of H₂O/AcOH (99:1) (solvent A) and MeOH (solvent B) starting with 35-80% of B (20 min), 80-92% of B (20-25 min), maintaining at 92% for 8 min, with a flow rate of 1 mL/min. Data acquisition was

accomplished with the Chromera® software version 3.4.1. Compounds were further analyzed by UV spectroscopy with the shift reagents AlCl_3 and AlCl_3/HCl .¹⁰

3-*O*-methylquercetin (**1**): yellow oil; UV (MeOH): λ_{max} (log ϵ): 245, 300, 355; (MeOH + AlCl_3): λ_{max} (log ϵ): 235, 265, 440; (MeOH + AlCl_3/HCl): λ_{max} (log ϵ): 230, 265, 405; ^1H NMR (500.00 MHz, CD_3OD): δ (ppm) = 6.21 (s, 1H, H-6), 6.40 (s, 1H, H-8), 7.65 (s, 1H, H-2'), 6.93 (d, J = 8.3 Hz, 1H, H-8'), 7.56 (d, J = 8.3 Hz, 1H, H-6'), 3.81 (s, 3H, OCH_3 -3); MS: m/z 317 $[\text{M}+\text{H}]^+$.

3,4'-*O*-dimethylquercetin (**2**): yellow oil; UV (MeOH): λ_{max} (log ϵ): 255, 295, 355; (MeOH + AlCl_3): λ_{max} (log ϵ): 235, 355, 415; (MeOH + AlCl_3/HCl): λ_{max} (log ϵ): 225, 270, 405; ^1H NMR (500.00 MHz, CD_3OD): δ (ppm) = 6.23 (d, J = 2.1 Hz, 1H, H-6), 6.44 (m, 1H, H-8), 7.75 (d, J = 2.1 Hz, 1H, H-2'), 6.98 (d, J = 8.5 Hz, 1H, H-5') e 7.67 (dd, J = 2.1, 8.5 Hz, 1H, H-6'), 3.83 (s, 3H, OCH_3 -3), 3.97 (s, 3H, OCH_3 -4'); ^{13}C NMR (125.0 MHz, CD_3OD): δ (ppm) = 56.4 (CH_3 , OMe -4'), 60.4 (CH_3 , OCH_3 -3), 94.7 (CH, C-8), 99.7 (CH, C-6), 112.8 (CH, C-2'), 116.3 (CH, C-5'), 138.2 (OCH_3 -3), 147.6 (OCH_3 -4'); MS: m/z 331 $[\text{M}+\text{H}]^+$.

3,7-*O*-dimethylquercetin (**3**): yellow oil; UV (MeOH): λ_{max} (log ϵ): 240, 300, 370; (MeOH + AlCl_3): λ_{max} (log ϵ): 235, 440; (MeOH + AlCl_3/HCl): λ_{max} (log ϵ): 235, 355, 410; ^1H NMR (500.00 MHz, CD_3OD): δ (ppm) = 8.6 (brs, OH-5), 6.35 (d, J = 2.2, 1H, H-6), 6.61 (d, J = 2.2, 1H, H-8), 7.65 (d, J = 2.2, 1H, H-2'), 4.83 (brs, OH-3'-4'), 6.91 (d, J = 8.5 Hz, H-5'), 7.56 (dd, J = 2.2, 8.5 Hz, H-6'), 3.80 (s, 3H, OCH_3 -3), 3.89 (s, 3H, OCH_3 -7); ^{13}C NMR (125.0 MHz, CD_3OD): δ (ppm) = 56.2 (CH_3 , OCH_3 -7), 60.2 (CH_3 , OCH_3 -3), 92.7 (CH, C-8), 98.6 (CH, C-6), 105.3 (C, C-10), 116.1 (CH, C-5'), 116.2 (CH, C-2'), 121.0 (C, C-1'), 122.1 (CH, C-6'), 138.3 (OCH_3 -3), 145.1 (C, OH-3'), 148.7 (C, OH-4'), 156.9 (C, C-2), 161.4 (C, OH-5), 165.9 (OCH_3 -7); MS: m/z 331 $[\text{M}+\text{H}]^+$.

2.5. Cytotoxic activity assay

Crude methanol extract (ME) was evaluated for *in vitro* cytotoxicity against three human tumor cell lines, melanoma (SK-Mel 28), gastric ascites (AGP-01) and breast carcinoma (MCF-7), and one non-tumor cell line, fetal lung fibroblast (MRC5), using the Alamar Blue (AB) assay.¹¹ Doxorubicin was used as positive control. The concentration of samples resulting in 50% growth inhibition (IC_{50}) was calculated for each cell line in GraphPad Prism® 5.0.

3. Results and discussion

Three flavonoids, 3-*O*-methylquercetin (**1**), 3,4'-*O*-dimethylquercetin (**2**) and 3,7-*O*-dimethylquercetin (**3**) were isolated from the methanolic extract of the bark and wood of *P. olacoides* (Fig. 1) for the first time. The structures of compounds are supported by 1D (^1H and ^{13}C) and 2D (COSY, HSQC and HMBC) NMR experiments, UV and LC-ESI-MS analysis, and are in agreement with those reported in the literature.¹² UV shifts data confirmed the position of the free hydroxyl groups. These confirmations were possible once the use of the shift reagents AlCl_3 and AlCl_3/HCl permits differentiation of the formation of acid-stable complexes between hydroxyls and neighboring ketones, and acid-labile complexes with *ortho*-dihydroxyl groups. Thus, the bathochromic shift caused by AlCl_3/HCl on band 1 of the spectra of **1-3** is characteristic of a free hydroxyl group in carbon atom C-5, and the absence of oxygen atom at carbon C-6 along with the bathochromic shift caused by AlCl_3 on band 1 of the spectra of **1** and **3**, is characteristic of *ortho*-dihydroxyl groups on B-ring.¹⁰

Crude methanol extract was evaluated *in vitro* for its cytotoxic activity against gastric ascites (AGP-01), breast (MCF-7) and melanoma (SK-Mel-28) cancer cells. The crude extract presented a moderate cytotoxic effect against MCF-7, with IC_{50} of 45.16 $\mu\text{g}/\text{mL}$, when compared with doxorubicin (positive control). In addition, the extract did

not display cytotoxicity against MRC-5 (non-tumor human fibroblast cells).

Although we have not studied the cytotoxic activity of the isolated flavonoids, literature reports that these quercetin derivatives exhibit a variety of biological activities, including antiproliferative and

antioxidant properties.¹³ For instance, Talib and co-workers^{13b} described the antiproliferative activity of **1** against MCF-7 cells with an IC₅₀ value of 11.23 µg/mL. Therefore, the compounds described in this paper may be important for the cytotoxic activity against MCF-7 cancer cells.




Figure 1. Structures of flavonoids **1-3** isolated from the methanolic extract of wood and bark of *Ptychopetalum olacoides* Benth

4. Conclusion

This is the first report on isolation of flavonoids and cytotoxic activity for the genus *Ptychopetalum*. Bioguided assays should be further performed in order to confirm that these flavonoids are the main active compounds involved in the cytotoxic activity.

Acknowledgements

The authors gratefully thank the financial support of FAPERJ for this study, the research stipends from CAPES to KDBD and, from CNPq, to ALM, ALV, RCM, RAE and TRAV.

References

¹ Silva, R. A. D., Plantas medicinais Brasileiras: estudo botânico e farmacognóstico - Muirapuama. *Revista Brasileira de Medicina e Pharmacia* **1925**, *1*, 37.

² Rossi, L., Olacaceae. Em *Lista de espécies da Flora do Brasil. Jardim Botânico do Rio de Janeiro*, Instituto de Pesquisas Jardim Botânico do Rio de Janeiro: Rio de Janeiro, Brazil, 2014; Vol. 2, pp 1339. [Link]

³ Siqueira, I. R.; Fochesatto, C. n.; Silva, A. L.; Nunes, D. S.; Battastini, A. M.; Netto, C. A.; Elisabetsky, E., *Ptychopetalum olacoides*, a traditional Amazonian "nerve tonic", possesses anticholinesterase activity. *Pharmacology Biochemistry and Behavior* **2003**, *75*, 645. [CrossRef] [PubMed]

⁴ Antunes, E.; Gordo, W. M.; Oliveira, J. F.; Teixeira, C. E.; Hyslop, S.; De Nucci, G., The relaxation of isolated rabbit corpus cavernosum by the herbal medicine Catuama® and its constituents. *Phytotherapy Research* **2001**, *15*, 416. [CrossRef] [PubMed]

⁵ Montrucchio, D. P.; Migeuel, O. G.; Miguel, M. D.; Monache, F. D.; Carvalho, J. L. S., Chemical compounds and antimicrobial activity of *Ptychopetalum olacoides* Benth. *Visão Acadêmica* **2005**, *6*, 48. [CrossRef]

⁶ a) Tang, W.; Hioki, H.; Harada, K.; Kubo, M.; Fukuyama, Y., Clerodane diterpenoids with NGF-potentiating activity from *Ptychopetalum olacoides*. *Journal of Natural Products* **2008**, *71*, 1760; [CrossRef]

- [PubMed] b) Tang, W.; Kubo, M.; Harada, K.; Hioki, H.; Fukuyama, Y., Novel NGF-potentiating diterpenoids from a Brazilian medicinal plant, *Ptychopetalum olacoides*. *Bioorganic & Medicinal Chemistry Letters* **2009**, *19*, 882; [CrossRef] [PubMed] c) Tang, W.; Harada, K.; Kubo, M.; Hioki, H.; Fukuyama, Y., Eight new clerodane diterpenoids from the bark of *Ptychopetalum olacoides*. *Natural Product Communications* **2011**, *6*, 327. [PubMed]
- ⁷ Colombo, R.; Batista, A. N. d. L.; Bomfim, G. C. C.; Burgos, R. C. R.; Cavalheiro, A. J.; Bolzani, V. S.; Silva, D. H. S.; Reimberg, M. C. H., Validated high-performance liquid chromatographic method for the standardisation of *Ptychopetalum olacoides* Benth., Olacaceae, commercial extracts. *Brazilian Journal of Pharmacognosy* **2010**, *20*, 781. [CrossRef]
- ⁸ Rolim, A.; Maciel, C. P. M.; Kaneko, T. M.; Consiglieri, V. O.; Salgado-Santos, I. M. N.; Velasco, M. V. R., Validation assay for total flavonoids, as rutin equivalents, from *Trichilia catigua* Adr. Juss (Meliaceae) and *Ptychopetalum olacoides* Bentham (Olacaceae) commercial extract. *Journal of AOAC International* **2005**, *88*, 1015. [PubMed]
- ⁹ Picerno, P.; Mencherini, T.; Rastrelli, L.; Piccinelli, A.; Aquino, R., Isoprenoid glycosides from *Liriosma ovata*. *Journal of Natural Products* **2008**, *71*, 265. [CrossRef] [PubMed]
- ¹⁰ Markham, U. R., Techniques of flavonoid identification. Em *Biol. Plantarum*, Challice, J., Ed. Academic Press: London, New York, 1982; Vol. 26, pp 36. [CrossRef]
- ¹¹ Nakayama, G. R.; Caton, M. C.; Nova, M. P.; Parandoosh, Z., Assessment of the Alamar Blue assay for cellular growth and viability *in vitro*. *Journal of Immunological Methods* **1997**, *204* (2), 205. [CrossRef] [PubMed]
- ¹² a) Kwon, Y. S.; Kim, C. M., Antioxidant constituents from the stem of *Sorghum bicolor*. *Archives of Pharmacal Research* **2003**, *26*, 535; [CrossRef] [PubMed] b) Costa, F. J.; Bandeira, P. N.; Albuquerque, M. R. J. R.; Pessoa, O. D. L.; Silveira, E. R.; Braz-Filho, R., Constituintes químicos de *Vernonia chalybaea* mart. *Química Nova* **2008**, *31*, 1691; [CrossRef] c) Rashed, K.; Sahuc, M.-E.; Deloison, G.; Calland, N.; Brodin, P.; Rouillé, Y.; Séron, K., Potent antiviral activity of *Solanum rantonnetii* and the isolated compounds against hepatitis C virus *in vitro*. *Journal of Functional Foods* **2014**, *11*, 185. [CrossRef]
- ¹³ a) Torres, F.; Quintana, J.; Estévez, F., 5,7,3'-trihydroxy-3,4'-dimethoxyflavone-induced cell death in human Leukemia cells is dependent on caspases and activates the MAPK pathway. *Molecular Carcinogenesis* **2010**, *49*, 464; [CrossRef] [PubMed] b) Talib, W. H.; Zarga, M. H. A.; Mahasneh, A. M., Antiproliferative, antimicrobial and apoptosis inducing effects of compounds isolated from *Inula viscosa*. *Molecules* **2012**, *17*, 3291; [CrossRef] [PubMed] c) Santos, K. P.; Motta, L. B.; Santos, D. Y. A. C.; Salatino, M. L. F.; Salatino, A.; Ferreira, M. J. P.; Lago, J. H. G.; Ruiz, A. L. T. G.; Carvalho, J. E.; Furlan, C. M., Antiproliferative activity of flavonoids from *Croton sphaerogynus* Baill. (Euphorbiaceae). *BioMed Research International* **2015**, *2015*, 7; [CrossRef] [PubMed] d) Shi, Z.-H.; Li, N.-G.; Tang, Y.-P.; Shi, Q.-P.; Zhang, W.; Zhang, P.-X.; Dong, Z.-X.; Li, W.; Zhang, X.; Fu, H.-A.; Duan, J.-A., Synthesis, biological evaluation and SAR analysis of *O*-alkylated analogs of quercetin for anticancer. *Bioorganic & Medicinal Chemistry Letters* **2014**, *24*, 4424; [CrossRef] [PubMed] e) Vargas, F. d. S.; Almeida, P. D. O.; Boleti, A. P. A.; Pereira, M. M.; Souza, T. P.; Vasconcellos, M. C.; Nunez, C. V.; Pohlit, A. M.; Lima, E. S., Antioxidant activity and peroxidase inhibition of Amazonian plants extracts traditionally used as anti-inflammatory. *BMC Complementary and Alternative Medicine* **2016**, *16*, 83. [CrossRef] [PubMed]