

Artigo

Chemical Constituents of Non-Polar Fractions Obtained from *Cnidoscolus quercifolius* Pohl (Euphorbiaceae), a Medicinal Plant Native from the Brazilian Caatinga Biome

Oliveira-Junior, R. G.; Ferraz, C. A. A.; Oliveira, A. P.; Alencar-Filho, J. M. T.; Araújo, E. C. C.; Nunes, X. P.; Picot, L.; Braz-Filho, R.; Rolim, L. A.; Almeida, J. R. G. S.*

Rev. Virtual Quim., 2019, 11 (2), no prelo. Data de publicação na Web: 8 de abril de 2019

<http://rvq.sbq.org.br>

Constituintes Químicos de Frações Não-Polares Obtidas de *Cnidoscolus quercifolius* Pohl (Euphorbiaceae), uma Planta Medicinal Nativa do Bioma Caatinga Brasileiro

Resumo: *Cnidoscolus quercifolius* Pohl é uma planta medicinal do Brasil, nativa do bioma Caatinga, que apresenta diversas propriedades farmacológicas. Nesse estudo, nós descrevemos a identificação de compostos de frações não-polares obtidas das folhas (Hex-Le) e cascas do caule (Hex-Sb) da espécie. As análises químicas foram realizadas por CG-EM. Os constituintes químicos foram identificados com base nos dados espectrais obtidos e por comparação com os dados da literatura. O perfil de fragmentação dos principais compostos identificados também foi proposto e é igualmente apresentado nesse trabalho. Hidrocarbonetos (46,84 %), triterpenos (32,60 %) e derivados de carotenoides (0,83 %) foram considerados os constituintes majoritários de Hex-Le. Em contraste, hidrocarbonetos (41,89 %), diterpenos (12,83 %) e triterpenos (7,0 %) foram encontrados em Hex-Sb. Quatro diterpenos (deidroabietano, sandaracopimaradieno, kaur-16-eno e 13-metil-17-norkaur-15-eno) e um triterpeno (diplopteno) são relatados pela primeira vez no gênero *Cnidoscolus*.

Palavras-chave: Plantas medicinais; Caatinga; terpenoides; *Cnidoscolus*.

Abstract

Cnidoscolus quercifolius Pohl is a Brazilian medicinal plant from the Caatinga biome which possesses several pharmacological properties. In this paper we describe the identification of compounds from non-polar fractions of the leaves (Hex-Le) and stem-barks (Hex-Sb) from *C. quercifolius*. Chemical analyses were performed by GC-MS approach. Chemical constituents were identified based on spectral data obtained and by comparison with literature data. Fragmentation profile of the main compounds was proposed and presented in this paper. Hydrocarbons (46.84 %), triterpenes (32.60 %) and carotenoid derivatives (0.83 %) were considered the major constituents of Hex-Le. In contrast, hydrocarbons (41.89 %), diterpenes (12.83 %) and triterpenes (7.0 %) were found in Hex-Sb. Four diterpenes (dehydroabietane, sandaracopimaradiene, kaur-16-ene and 13-methyl-17-norkaur-15-ene) and one triterpene (diploptene) are reported for the first time in the *Cnidoscolus* genus.

Keywords: Medicinal plants; Caatinga; terpenoids; *Cnidoscolus*.

* Universidade Federal do Vale do São Francisco, Núcleo de Estudos e Pesquisas de Plantas Medicinais (NEPLAME), CEP 56304-205, Petrolina-PE, Brazil.



jackson.guedes@univasf.edu.br

DOI:

Chemical Constituents of Non-Polar Fractions Obtained from *Cnidoscolus quercifolius* Pohl (Euphorbiaceae), a Medicinal Plant Native from the Brazilian Caatinga Biome

Raimundo G. de Oliveira-Júnior,^a Christiane Adrielly A. Ferraz,^a Ana Paula de Oliveira,^a José Marcos T. de Alencar-Filho,^a Edigênia C. C. Araújo,^a Xirley P. Nunes,^a Laurent Picot,^b Raimundo Braz-Filho,^c Larissa A. Rolim,^a Jackson Roberto G. da S. Almeida^{a,*}

^a Universidade Federal do Vale do São Francisco, Núcleo de Estudos e Pesquisas de Plantas Medicinais (NEPLAME), CEP 56304-205, Petrolina-PE, Brasil.

^b Université de La Rochelle, Faculté des Sciences et Technologies, UMRi CNRS 7266 LIENSs, La Rochelle, France.

^c Universidade Estadual do Norte Fluminense, Departamento de Química de Produtos Naturais, CEP 28035-200, Rio de Janeiro-RJ, Brasil.

* jackson.guedes@univasf.edu.br

Recebido em 6 de março de 2019. Aceito para publicação em 6 de março de 2019

1. Introduction

2. Experimental

2.1. Plant material

2.2. Chemicals

2.3. Extraction and fractionation

2.4. Gas Chromatography-Mass Spectrometry (GC-MS) analysis

2.5. Identification of compounds

3. Results and Discussion

4. Conclusions

1. Introduction

Caatinga is an exclusively Brazilian biome, which occurs predominantly in the Northeastern Region and comprises approximately 10 % of the Brazilian territory. This biome usually presents dry vegetation, characterized by semiarid climate, low and

irregular rainfall and fertile soil.^{1,2} Despite the adverse climatic conditions, Caatinga flora is very diverse and includes numerous species used as food source or in folk medicine. In a previous study, the use of 385 Caatinga species for medicinal purposes was verified. These species were distributed in about 265 genera and 91 different families.³ Although the Caatinga biome presents several species

with therapeutic potential, chemical and pharmacological investigations involving these plants are still scarce,^{4,5} which makes this biome one of the least studied in the world.

Cnidoscolus quercifolius (syn. *C. phyllacanthus* (Mull. Arg.) Pax & L. Hoffm.) is a medicinal plant native to Caatinga, popularly known as favela, faveleira or urtiga-branca. In folk medicine, its leaves and stem barks are used to treat hemorrhoids, renal problems, ophthalmic diseases, injury, skin problems, urinary tract infection and inflammatory processes.³ In fact, pharmacological investigations have demonstrated that extracts or isolated compounds from *C. quercifolius* have some therapeutic properties, such as antinociceptive,⁶ anti-inflammatory,⁷ antioxidant,⁸⁻¹⁰ cytotoxic¹¹ and antimicrobial⁹ activities.

The therapeutic properties of *C. quercifolius* are justified by the presence of bioactive secondary metabolites, mainly terpenoids. Phytochemical studies have resulted in the isolation of new tricyclic benzocyclohepten diterpenes, called favelins, a reference to the popular name of the species.¹²⁻¹⁴ Endo et al.¹⁵ also reported the isolation of favelanone and neofavelanone, new cyclopropane and cyclobutene tetracyclic derivatives, respectively. To date, these diterpenes have been reported exclusively in *C. quercifolius*, indicating that these compounds can be considered its chemotaxonomic markers. However, its chemical composition is still poorly known. In this paper, we describe the identification of non-polar compounds from leaves and stem-barks fractions of *C. quercifolius*, including compounds reported for the first time in the genus.

2. Experimental

2.1. Plant material

Leaves and stem-barks of *Cnidoscolus quercifolius* Pohl were collected in Petrolina, in a Caatinga area (coordinates 09° 03' 55.30" S and 40° 20' 06.90" W), State of Pernambuco, Brazil, in February 2013. A voucher specimen (n° 19202) was deposited at the Herbário Vale do São Francisco (HVASF), of the Universidade Federal do Vale do São Francisco (UNIVASF). All procedures for access to genetic patrimony and associated traditional knowledge were carried out and the project was registered in SisGen (Register #A65F584).

2.2. Chemicals

Homologous series of *n*-alkanes (C₉H₂₀ – C₄₀H₈₂) were purchased from Merck® (Germany). All solvents (ethanol, hexane, chloroform, ethyl acetate and methanol) were purchased from Synth® (Brazil).

2.3. Extraction and fractionation

The dried and pulverized leaves (482 g) and stem-barks (1,481 g) of *C. quercifolius* were separately subjected to maceration with 95 % ethanol for 72 h. After, the solution was filtered and concentrated under reduced pressure on a rotatory evaporator at 50 °C, producing 63 and 392 g of ethanol extract from leaves (EtOH-Le, 13.07 %) and stem-barks (EtOH-Sb, 26.46 %). Subsequently, an aliquot of EtOH-Le (15 g) and EtOH-Sb (50 g) was fractionated by vacuum liquid chromatography (VLC), using silica gel 60 as stationary phase. Hexane (Hex), chloroform (CHCl₃), ethyl acetate (AcOEt) and methanol (MeOH) were used as mobile phase in ascending order of polarity, resulting in the respective fractions, as shown in figure 1. Hexane fractions (Hex-Le and Hex-Sb) were chosen for GC-MS analysis.




Figure 1. Fractionation of EtOH-Le and EtOH-Sb by vacuum liquid chromatography (VLC). Hexane fractions (Hex-Le and Hex-Sb) were selected for GC-MS analysis

2.4. Gas Chromatography - Mass Spectrometry (GC-MS) analysis

Qualitative and quantitative determination of the chemical constituents present in Hex-Le and Hex-Sb was performed by GC-MS using a Shimadzu® gas chromatograph (QP-2010) interfaced with a mass spectrometer, employing the following chromatographic conditions: Phenomenex® ZB-5MS Zebron column (30.0 m x 0.25 mm x 0.25 mm); helium (99.999 %) carrier gas at a constant flow of 1.1 ml/min; 1 µl injection volume; injector split ratio of 1:40; injector temperature 240 °C; electron impact mode at 70 eV; ion source temperature 280 °C. The oven temperature was programmed at 100 °C (isothermal for 5 min), with an increase of 10 °C/min to 250 °C (isothermal for 5 min) and 10 °C/min to 280 °C (isothermal for 5 min). A mixture of linear hydrocarbons (C₉H₂₀ – C₄₀H₈₂) was injected under the same experimental conditions.

2.5. Identification of compounds

Constituents were identified by comparison of their mass spectra with those of authentic compounds or with reference spectra in the computer library (Wiley7lib and NIST08lib). For some identified

secondary metabolites (triterpenes, diterpenes and carotenoid derivatives), the fragmentation profile was proposed based on the peaks observed in the mass spectrum.

3. Results and Discussion

GC-MS chromatogram of Hex-Le revealed the presence of 49 peaks, of which 30 were identified, corresponding to 80.27 % of its total chemical composition (Figure 2A). Among the identified compounds, 0.83 % were carotenoid derivatives, 46.84 % were long chain hydrocarbons or derivatives, and 32.60 % corresponded to squalene, a triterpene considered the major constituent of the sample. Regarding Hex-Sb, the chromatogram showed 74 peaks (Figure 2B), of which 46 were identified, corresponding to 61.72 % of its chemical composition. Most of the compounds were long chain hydrocarbons or derivatives (41.89 %), followed by diterpenes (12.83 %) and triterpenes (7.0 %). Carotenoid derivatives were not identified in this sample. The major chemical constituents of Hex-Sb were hexadecane (7.48 %) and sandaracopyramadiene (7.53 %). All identified compounds are described in Table 1.

Two carotenoid derivatives were identified in Hex-Le, β -ionone (0.26 %) and dihydroactinidiolide (0.57 %) (Figure 3). These constituents are considered products of carotenoid degradation through the action of carotenoid dioxygenase enzymes.^{16,17} In fact, carotenoid degradation products indicate that the plant is in an oxidative stress condition, frequently associated with the presence of singlet oxygen ($O_2^{\cdot}$).^{18,19} A summary of the biosynthesis as well as the proposed fragmentation profile for β -ionone and dihydroactinidiolide are shown in figure 4. Initially, an α -carotene molecule is used by the carotenoid dioxygenase enzyme as substrate, producing α -ionone and 10'-apo- β -10'-carotenal, which is then converted to β -ionone (m/z 192). In the mass spectra, β -ionone loses a methyl group by a cleavage adjacent to the annular double bond, yielding a new fragment (m/z 177), compatible with the base peak (Figure 3). In an oxidative stress situation, β -ionone molecule is readily oxidized to 5,6-epoxy- β -ionone and, after a rearrangement step, leads to the formation of dihydroactinidiolide (m/z 180).^{16,17} Rupture of the lactone ring of this molecule releases a

stable fragment (m/z 111), compatible with the base peak observed in the mass spectra (Figure 3).

The triterpene squalene was found in lower amount in Hex-Sb (4.65 %). However, other triterpenes were identified in this sample (diploptene 0.20 %, lupeol 2.15 %) and its mass spectra are shown in Figure 5. Squalene is considered the precursor of diploptene and lupeol, as shown in Figure 6. In its mass spectrum, m/z 69 (base peak) and 81 were the most stable fragments, already described in the literature as characteristic fragments of this molecule.²⁰ For lupeol and diploptene, spectral data contributed significantly to their characterization through detection of molecular ions (m/z 426 and m/z 410, respectively) and base peaks (m/z 191 and m/z 218, respectively).^{21,22} Besides these, other characteristic fragments are shown in figure 7. Paula et al.¹⁴ have reported the presence of lupeol and derivatives in *C. quercifolius* extracts. However, there are no reports of the presence of diploptene in *Cnidioscolus* species to date.




Figure 2. GC-MS chromatograms obtained for Hex-Le (A) and Hex-Sb (B)

Table 1. Chemical constituents of non-polar fractions from leaves (Hex-Le) and stem-bark (Hex-Sb) of *Cnidocolus quercifolius*. Compounds were identified by CG-MS analysis

Phytochemical	Compound	Hex-Le		Hex-Sb	
		RT (min)	Content (%)	RT (min)	Content (%)
Carotenoid derivatives	β -ionone	22.84	0.26	ND	ND
	Dihydroactinidiolide	24.41	0.57	ND	ND
Identified (%)		-	0.83	-	ND
Hydrocarbons and derivatives	1-tridecene	ND	ND	19.45	0.28
	Tetradecane	19.80	0.56	19.82	3.00
	2,3,7-trimethyl-decane	ND	ND	22.27	0.12
	2-methyl-tetradecane	ND	ND	22.37	0.13
	3-methyl-tetradecane	ND	ND	22.65	0.13
	2-methyl-hexadecan-1-ol	ND	ND	23.48	0.21
	Pentadecane	23.81	0.99	23.82	1.99
	5-methyl-pentadecane	ND	ND	25.83	0.29
	2-methyl-pentadecane	ND	ND	26.29	0.24
	3-methyl-pentadecane	26.53	0.18	26.55	0.59
	3-hexadecene	ND	ND	26.82	0.26
	1-hexadecene	ND	ND	27.37	1.85
	Hexadecane	27.67	2.93	27.70	7.48
	3-hexyl-1,1,2-trimethyl-cyclobutane	ND	ND	27.84	0.15
	2,6,10-trimethyl-pentadecane	29.45	0.56	29.45	0.15
	1-cyclohexyl-decane	29.69	0.17	ND	ND
	1-decyl-cyclopentane	ND	ND	29.70	0.19

2-methyl-hexadecane	ND	ND	30.03	0.17
2-methyl-heptadecane	ND	ND	30.29	0.11
2-phenyl-dodecane	ND	ND	31.07	0.26
Heptadecane	31.36	1.79	31.37	3.01
2,6,10,14-tetramethyl-pentadecane	31.52	1.40	31.51	0.50
7-methyl-hexadecane	ND	ND	32.94	0.75
4-ethyl-heptadecane	33.15	0.26	ND	ND
3-methyl-heptadecane	33.85	0.24	33.87	0.46
1-octadecene	ND	ND	34.61	1.88
Octadecane	ND	ND	34.61	1.88
2,6,10,14-tetramethyl-hexadecane	35.11	1.15	ND	ND
4-cyclohexyl-tridecane	37.00	0.18	ND	ND
Hexadecan-1-ol	37.48	0.54	ND	ND
Nonadecane	38.22	1.69	38.24	0.91
Octasane	40.26	0.18	ND	ND
3-methyl-nonadecane	40.52	0.19	ND	ND
2-methyl-eicosane	ND	ND	40.53	0.36
1-tricosene	41.20	0.21	47.23	0.83
1-nonadecene	ND	ND	41.21	1.06
Eicosane	41.43	1.85	41.65	0.68
Octadecan-1-ol	43.91	0.17	ND	ND
Heneicosan-1-ol	ND	ND	44.28	0.32
Heneicosane	44.48	1.46	44.49	2.36
7-hexyl-eicosane	ND	ND	46.60	0.19

	Docosane	47.41	1.34	47.43	3.22
	7-hexyl-docosane	ND	ND	52.19	0.14
	Pentacosane	60.43	5.69	55.35	2.96
	Nonacosane	55.52	3.09	ND	ND
	Hexacosane	62.83	1.79	58.02	1.09
	Heptacosan-1-ol	67.06	0.55	60.31	0.60
	Tetracontane	69.27	9.59	64.99	1.09
	Tetratetracontane	73.27	8.09	ND	ND
Identified (%)		-	46.84	-	41.89
<i>Diterpenes</i>	Sandaracopimaradiene	ND	ND	39.63	7.53
	13-methyl-17-norkaur-15-ene	ND	ND	41.65	0.68
	Kaur-16-ene	ND	ND	42.66	4.38
	Dehydroabietane	ND	ND	42.89	0.24
Identified (%)		-	ND	-	12.83
<i>Triterpenes</i>	Squalene	63.09	32.60	63.03	4.65
	Lupeol	ND	ND	69.18	2.15
	Diploptene	ND	ND	73.35	0.20
Identified (%)		-	32.60	-	7.00
Total identified (%)			80.27		61.72

RT: retention time. ND: compound not detected in sample.




Figure 3. Mass spectrum obtained for β -ionone and dihydroactinidiolide identified in Hex-Le




Figure 4. Biosynthesis and proposed fragmentation of carotenoid derivatives β -ionone and dihydroactinidiolide




Figure 5. Mass spectra obtained for the triterpenes identified in Hex-Sb




Figure 6. Simplified biosynthetic route for diploptene and lupeol, and proposed fragmentation profile for squalene




Figure 7. Proposed fragmentation of lupeol and diploptene, identified in Hex-Sb

GC-MS analysis also showed abietane (dehydroabietane 0.24 %), pimarane (sandaracopimaradiene 7.53 %) and kaurene (kaur-16-ene 4.38 %, 13-methyl-17-norkaur-15-ene 0.68 %) diterpenes in Hex-Sb, reported for the first time in *Cnidoscopus*. Figure 8 shows the mass spectra of

sandaracopimaradiene as well as the proposed fragmentation profile for the molecule. An abundant fragment (m/z 257) was registered after loss of the methyl group, while the base peak (m/z 137) was originated after a B-ring cleavage, as shown in the figure.




Figure 8. Mass spectrum and fragmentation proposed for sandaracopimaradiene, identified in Hex-Sb

Figure 9 shows dehydroabietane mass spectra and its fragmentation profile. Initially, the molecule loses a methyl group, yielding a stable fragment, compatible with the base peak (m/z 255). From that fragment, one

isopropyl group is removed, affording m/z 213 fragment. In addition, A-ring cleavage has been proposed, yielding the fragments m/z 185 and m/z 159.




Figure 9. Mass spectrum and fragmentation proposed for dehydroabietane, identified in Hex-Sb

In the mass spectra of kaur-16-ene, it was observed a molecular ion peak (m/z 272), compatible with the molecular formula $C_{20}H_{32}$. After ionization, this diterpene also loses a methyl group, providing a stable fragment (m/z 257). Subsequently, C-ring undergoes a retro Diels-Alder cleavage, yielding m/z 229 fragment. Hydrogen rearrangements were proposed, resulting in new fragments (m/z 187, 147, 213 and 133), as shown in the figure 10. In addition, D-ring hydrogen rearrangements led to the formation of a stable fragment, attributed to the tropylium ion (m/z 91), compatible with the base peak found in the mass spectrum.

For 13-methyl-17-norkaur-15-ene, fragmentation begins with the loss of C-13

methyl, generating a relatively stable fragment (m/z 257). This fragment undergoes a series of rearrangements, resulting in m/z 229 and m/z 119 peaks. The m/z 257 fragment undergoes a B-ring cleavage and then rearrangement of hydrogens, releasing a new fragment at m/z 134. This fragment loses a methyl group, resulting in the m/z 119 peak. Formation of the fragment corresponding to the base peak of the spectrum (m/z 79) occurs by cleavage of the last fragment recorded at m/z 119, as shown in figure 11. The complete fragmentation profile of the diterpenes identified in Hex-Sb is shown in the figures 8-11.




Figure 10. Mass spectrum and fragmentation proposed for kaur-16-ene, identified in Hex-Sb




Figure 11. Mass spectrum and fragmentation proposed for 13-methyl-17-norkaur-15-ene, identified in Hex-Sb

4. Conclusions

In summary, this study reported a chemical investigation of non-polar fractions obtained from leaves and stem-barks of *C. quercifolius*. We describe for the first time the identification of diterpenes dehydroabietane, sandaracopimaradiene, kaur-16-ene and 13-methyl-17-norkaur-15-ene, in a *Cnidioscolus* species. Indicators of oxidative stress were

also identified (β -ionone and dihydroactinidiolide), in addition to triterpenes (lupeol and diploptene) commonly found in plant species. In this sense, our investigation has contributed to the chemical knowledge of *C. quercifolius*, a Brazilian medicinal plant from Caatinga biome.

Acknowledgments

The authors thanks to FACEPE (Fundação de Amparo à Ciência e Tecnologia do Estado de Pernambuco) for financial Support.

References

- ¹ Santos, K. G.; Dantas-Filho, A. N.; Leite, R. H. L.; Aroucha, E. M. M.; Santos, A. G.; Oliveira, T. A. Rheological and some physicochemical characteristics of selected floral honeys from plants of Caatinga. *Anais da Academia Brasileira de Ciências* **2014**, *86*, 981. [CrossRef]
- ² Albuquerque, U. P.; Araujo, E. L.; El-Deir, A. C. A.; Lima, A. L. A.; Souto, A.; Bezerra, B. M.; Ferraz, E. M. N.; Freire, E. M. X.; Sampaio, E. V. S. B.; Las-Casas, F. M. G.; Moura, G. J. B.; Pereira, G. A.; Melo, J. G. M.; Ramos, M. A.; Rodal, M. J. N.; Schiel, N.; Lyra-Neves, R. M.; Alves, R. R. N.; Azevedo-Junior, S. M.; Telino-Junior, W. R.; Severi, W. Caatinga Revisited: Ecology and Conservation of an Important Seasonal Dry Forest. *The Scientific World Journal* **2012**, *2012*, 1. [CrossRef] [PubMed]
- ³ Albuquerque, U. P.; Medeiros, P. M.; Almeida, A. L. S.; Monteiro, J. M.; Lins-Neto, E. M. F.; Gomes, J. M.; Santos, J. P. Medicinal plants of the Caatinga (semi-arid) vegetation of NE Brazil: a quantitative approach. *Journal of Ethnopharmacology* **2007**, *114*, 325. [CrossRef] [PubMed]
- ⁴ Menezes, R. S. C.; Sampaio, E. V. S. B.; Giongo, V.; Pérez-Marin, A. M. Biogeochemical cycling in terrestrial ecosystems of the Caatinga Biome. *Brazilian Journal of Biology* **2012**, *72*, 643. [CrossRef] [PubMed]
- ⁵ Albuquerque, U. P. Re-examining hypotheses concerning the use and knowledge of medicinal plants: a study in the Caatinga vegetation of NE Brazil. *Journal of Ethnobiology and Ethnomedicine* **2006**, *2*, 1. [CrossRef] [PubMed]
- ⁶ Gomes, L. M. C.; Lima-Saraiva, S. R. G.; Andrade, T. M. D.; Silva, J. C.; Diniz, T. C.; Barreto, V. N. S.; Mendes, R. L.; Quintans-Júnior, L. J.; Quintans, J. S. S.; Lima, J. T.; Almeida, J. R. G. S. Antinociceptive activity of the ethanolic extract from barks and leaves of *Cnidoscolus quercifolius* (Euphorbiaceae) in mice. *Journal of Young Pharmacists* **2014**, *6*, 64. [CrossRef]
- ⁷ Gomes, L. M. A.; Andrade, T. M. D.; Silva, J. C.; Lima, J. T.; Quintans-Júnior, L. J.; Almeida, J. R. G. S. Phytochemical screening and anti-inflammatory activity of *Cnidoscolus quercifolius* (Euphorbiaceae) in mice. *Pharmacognosy Research* **2014**, *6*, 345. [CrossRef] [PubMed]
- ⁸ Morais, N. R. L.; Oliveira-Neto, F. B.; Melo, A. R.; Bertini, L. M.; Silva, F. F. M.; Alves, L. A. Prospecção fitoquímica e avaliação do potencial antioxidante de *Cnidoscolus phyllacanthus* (müll. Arg.) Pax & k.hoffm. Oriundo de Apodi – RN. *Revista Brasileira de Plantas Medicinais* **2016**, *18*, 180. [CrossRef]
- ⁹ Paredes, P. F. M.; Vasconcelos, F. R.; Paim, R. T. T.; Marques, M. M. M.; Morais, S. M.; Lira, S. M.; Braquehais, I. D.; Vieira, I. G. P.; Mendes, F. N. P.; Guedes, M. I. F. Screening of Bioactivities and Toxicity of *Cnidoscolus quercifolius* Pohl. *Evidence-Based Complementary and Alternative Medicine* **2016**, *2016*, 1. [CrossRef]
- ¹⁰ Santos, K. A.; Aragão-Filho, O. P.; Aguiar, C. M.; Milinsk, M. C.; Sampaio, S. C.; Palú, F.; Silva, E. S. Chemical composition, antioxidant activity and thermal analysis of oil extracted from favela (*Cnidoscolus quercifolius*) seeds. *Industrial Crops and Products* **2017**, *97*, 368. [CrossRef]
- ¹¹ Peixoto-Sobrinho, T. J.; Tavares, E. A.; Castro, V. T. N. A.; Veras-Filho, J.; Militão, G. C. G.; Silva, T. G.; Amorim, E. L. C. Antiproliferative activity of species of the genus *Cnidoscolus* against HT-29, Hep-2 and NCI-H292 cells. *Molecular and Clinical Pharmacology* **2012**, *3*, 55.
- ¹² Endo, Y.; Ohta, T.; Nozoe, S. Favelines, novel tricyclic benzocycloheptenes with

cytotoxic activities from Brazilian plant, *Cnidoscopus phyllacanthus*. *Tetrahedron Letters* **1991**, 32, 3083. [[CrossRef](#)]

¹³ Lemos, T. L. G.; Silveira, E. R.; Oliveira, M. F.; Braz-Filho, R. Terpenoids from *Cnidoscopus phyllacanthus* Pax et Hoff. *Journal of the Brazilian Chemical Society* **1991**, 2, 105.

¹⁴ Paula, A. C.; Melo, K. M.; Silva, A. M.; Ferreira, D. A.; Monte, F. J. Q.; Santiago, G. M. P.; Lemos, T. L. G.; Braz-Filho, R.; Militão, G. C. G.; Silva, P. B. N.; Silva, T. G. Constituintes Químicos e Atividade Citotóxica de *Cnidoscopus phyllacanthus*. *Revista Virtual de Química* **2016**, 8, 231. [[CrossRef](#)]

¹⁵ Endo, Y.; Ohta, T.; Nozoe, S. Favelanone, a novel tetracyclic cyclopropane derivative from the Brazilian plant, *Cnidoscopus phyllacanthus*. *Tetrahedron Letters* **1991**, 32, 5555. [[CrossRef](#)]

¹⁶ Lima, K. S. C.; Lima, A. L. S.; Freitas, L. C.; Della-Modesta, R. C.; Godoy, R. L. O. Efeito de baixas doses de irradiação nos carotenoides majoritários em cenouras prontas para o consumo. *Ciência e Tecnologia de Alimentos* **2004**, 24, 183. [[CrossRef](#)]

¹⁷ Uenojo, M.; Maróstica-Junior, M. R.; Pastore, G. M. Carotenóides: propriedades, aplicações e biotransformação para formação de compostos de aroma. *Química Nova* **2007**, 30, 616. [[CrossRef](#)]

¹⁸ Ramel, F.; Birtic, S.; Ginies, C.; Soubigou-Taconnat, L.; Triantaphylidès, C.; Havaux, M. Carotenoid oxidation products are stress signals that mediate gene responses to singlet oxygen in plants. *Proceedings of the National Academy of Sciences* **2012**, 109, 5535. [[CrossRef](#)] [[PubMed](#)]

¹⁹ Havaux, M. Carotenoid oxidation products as stress signals in plants. *The Plant J*, 2013, 79, 597. [[CrossRef](#)] [[PubMed](#)]

²⁰ Oyugi, D. A.; Ayorinde, F. O.; Gugssa, A.; Allen, A.; Izevbigie, E. B.; Eribo, B.; Anderson, W. A. Biological activity and mass spectrometric Analysis of *Vernonia amygdalina* fractions. *Journal of Bioscience and Technology* **2011**, 2, 287.

²¹ Carvalho, T. C.; Polizeli, A. M.; Turatti, I. C.; Severiano, M. E.; Carvalho, C. E.; Ambrósio, S. R.; Crotti, A. E.; Figueiredo, U. S.; Vieira, P. C.; Furtado, N. A. Screening of Filamentous Fungi to Identify Biocatalysts for Lupeol Biotransformation. *Molecules* **2010**, 15, 6140. [[CrossRef](#)] [[PubMed](#)]

²² Bezerra, M. Z. B.; Campelo, P. A.; Machado, M. I. L.; Matos, F. J. A.; Braz-Filho, R. Constituintes químicos isolados de três espécies do gênero *Sclerolobium*. *Química Nova* **1994**, 17, 205. [[Link](#)]